

ORIGINAL INVESTIGATION

Open Access



Association of TyG index and TG/HDL-C ratio with arterial stiffness progression in a non-normotensive population

Zhiyuan Wu^{1,3†}, Di Zhou^{1†}, Yue Liu¹, Zhiwei Li¹, Jinqi Wang¹, Ze Han¹, Xinlei Miao¹, Xiangtong Liu¹, Xia Li², Wei Wang³, Xiuhua Guo^{1,2*} and Lixin Tao^{1,2*} 

Abstract

Background: Cross-sectional studies have reported that insulin resistance (IR) is associated with arterial stiffness. However, the relationship between IR and arterial stiffness progression remains unclear. This study aims to evaluate the association of triglyceride glucose (TyG) index and triglyceride to high-density lipoprotein cholesterol (TG/HDL-C) ratio with arterial stiffness progression in a non-normotensive population.

Methods: A total of 1895 prehypertensive (systolic pressure 120–139 mmHg or diastolic pressure 80–90 mmHg) or hypertensive (systolic pressure \geq 140 mmHg or diastolic pressure \geq 90 mmHg or using antihypertensive medication) participants were enrolled in 2013 and 2014, and followed until December 31, 2019. Arterial stiffness progression was measured by brachial-ankle pulse wave velocity (baPWV) change (absolute difference between baseline and last follow-up), baPWV change rate (change divided by following years), and baPWV slope (regression slope between examination year and baPWV).

Results: During a median follow-up of 4.71 years, we observed an increasing trend of baPWV in the population. There were linear and positive associations of the TyG index and TG/HDL-C ratio with the three baPWV parameters. The difference (95% CI) in baPWV change (cm/s) comparing participants in the highest quartile versus the lowest of TyG index and TG/HDL-C ratio were 129.5 (58.7–200.0) and 133.4 (52.0–214.9), respectively. Similarly, the evaluated baPWV change rates (cm/s/year) were 37.6 (15.3–60.0) and 43.5 (17.8–69.2), while the slopes of baPWV were 30.6 (9.3–51.8) and 33.5 (9.0–58.0). The observed association was stronger in the hypertensive population.

Conclusion: Our study indicates that the TyG index and TG/HDL-C ratio are significantly associated with arterial stiffness progression in hypertensive population, not in prehypertensive population.

Keywords: Insulin resistance, Arterial stiffness, Hypertension, Triglyceride glucose index, TG/HDL-C ratio, Cohort study

Background

Hypertension has caused a heavy economic burden worldwide, becoming a challenging public health issue. In 2010, 31.1% of adults around the world were reported to have hypertension [1]. Among Chinese adults aged 35–75 years, nearly half are diagnosed with hypertension, the incidence is still steadily increasing, and the onset age is becoming younger [2]. Among hypertensive patients, arterial stiffness is a common vascular complication and

*Correspondence: statguo@ccmu.edu.cn; taolixin@ccmu.edu.cn

[†]Zhiyuan Wu and Di Zhou are first co-authors and drafted the manuscript together

² Department of Mathematics and Statistics, La Trobe University, Melbourne, Australia

Full list of author information is available at the end of the article



is also an independent risk factor and predictor of other cardiovascular and cerebrovascular diseases [3, 4], such as coronary heart disease (CHD) and stroke. Therefore, it is of great importance to focus on the progression of arterial stiffness in the hypertensive population and identify the early related factors of arterial stiffness [5–7].

Disorders of glucose and lipid metabolism are a common pathophysiological feature accompanying patients with hypertension, while insulin resistance (IR) extensively participates in this biological process [8]. The hyperinsulinaemic-euglycaemic clamp is the gold standard for evaluating the status of IR [9]. However, this assessment process is expensive and complex and is not ideal for routine clinical monitoring. Recently, some novel and simple indicators have been reported to be reliable surrogate indexes of IR, such as the triglyceride-glucose (TyG) index and the triglyceride to high-density lipoprotein cholesterol (TG/HDL-C) ratio. Compared with the hyperinsulinaemic-euglycaemic clamp, the TyG index has a high sensitivity of 96.5% and a specificity of 85.0% for the diagnosis of IR [10], which has also been reported to be associated with diabetes in Chinese population [11]. Many studies have found that these surrogate indexes are independent risk factors for some cerebrocardiovascular diseases [12–14]. Moreover, IR-related indexes are associated with arterial stiffness [15–19]. However, these studies are all based on general populations. Recently, Li et al. [20] found that the TyG index is positively associated with brachial-ankle pulse wave velocity (baPWV) in hypertensive patients. However, this study is a cross-sectional study without follow-up. Moreover, the association between the TG/HDL-C ratio and




Fig. 1 Flow chart of the study population

was defined as SBP ≥ 140 mmHg or DBP ≥ 90 mmHg or use of any antihypertensive medication. Prehypertension status was defined as SBP of 120–139 mmHg or DBP of 80–89 mmHg. The mean arterial pressure (MAP) was calculated as $1/3 \times \text{SBP} + 2/3 \times \text{DBP}$. A higher vascular stiffness increases pulse pressure (PP), while PP could increase SBP and decrease DBP to a small extent.

Therefore, the MAP level was adjusted in the following analyses.

Blood samples were stored and measured in the central laboratory of Beijing Xiaotangshan Examination Center or Beijing Physical Examination Center using the Olympus Automatic Biochemical Analyser (Hitachi 747; Tokyo, Japan). Serum total cholesterol (TC), triglyceride (TG), high-density lipoprotein cholesterol (HDL-C), and low-density lipoprotein cholesterol (LDL-C) were measured with the enzymatic colour metric method. The estimated glomerular filtration rate (eGFR) was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI 2009) serum creatinine equation [23]. Fasting blood glucose (FBG) was defined as the glucose concentrations before breakfast after overnight fasting (no food, except drinking water, for at least 8–10 h), while two-hour postprandial blood glucose (PBG) was measured 2 h after the beginning of fixed

meals through finger blood tests. Diabetes was defined as FBG ≥ 7.0 mmol/L, PBG ≥ 11.1 mmol/L, glycated haemoglobin (HbA1c) $\geq 6.5\%$ or the use of any glucose-lowering medication according to the American Diabetes Association [24]. According to the Guidelines on Prevention and Treatment of Dyslipidaemia for Chinese Adults [25], dyslipidaemia was defined as TG ≥ 2.3 mmol/L, TC ≥ 6.2 mmol/L, LDL-C ≥ 4.1 mmol/L, HDL-C < 1.0 mmol/L, or any lipid-lowering medication. The TyG index was denoted as $\ln[\text{TG (mm/L)} \times \text{fasting glucose (mm/L)} / 2]$. The TG/HDL-C ratio was calculated as TG (mm/L) divided by HDL-C (mm/L).

Assessment of baPWV

baPWV is a simple, noninvasive, automatic and widely used method in clinical practice and large population-based studies. The baPWV was measured with an Omron Colin BP-203RPE III device (Omron Health Care, Kyoto, Japan). After more than 5 min of rest in the supine position, 4 cuffs were wrapped around the bilateral brachia and ankles and then connected to a plethysmographic sensor and oscillometric pressure sensor. Semiconductor pressure sensors were used to assess the transmission time between the initial rises in both the brachial and tibial artery waves to record the pressure waveform.

the distance between sampling points of baPWV was determined based on the height of the subjects. The time interval between the wave front of the brachial waveform and the ankle waveform was expressed as the time interval between the brachium and ankle (T_{ba}). The final baPWV was calculated as

$$\text{baPWV} = (L_a - L_b) / \Delta T_{ba} \quad (1)$$

L_a was the path length from the suprasternal notch to the ankle, and L_b was the path length from the suprasternal notch to the brachium using the following equations:

$$L_a = 0.8129 \times \text{height of the participant (in cm)} + 12.328 \quad (2)$$

$$L_b = 0.2195 \times \text{height of the participant (in cm)} - 2.0734 \quad (3)$$

The detailed process has been described in a previous study [26]. The maximum value of baPWV on the left and right sides was chosen as the final result of baPWV. The baPWV was measured at baseline and at the follow-up visit. The baPWV change was calculated as the baPWV value at baseline minus the baPWV value at the last visit during follow-up. The baPWV change rate was calculated as the value of baPWV change divided by the time distance between baseline and last visit. The slope of baPWV was defined as the linear regression slope with examination year as the independent variable and multiple baPWV measurements as the dependent variable.

Statistical analysis

Baseline characteristics are presented as the mean (standard deviation, SD), median [interquartile range, IQR] or number (percentage), as appropriate. Differences in baPWV change, baPWV change rate and slope of baPWV among groups were compared using the Kruskal–Wallis test.

Multivariate linear regression models were used to estimate the association of the TyG index and TG/HDL-C ratio with the absolute change, change rate and slope of baPWV. The TyG index and TG/HDL-C ratio were both analysed as continuous variables and categorized into quartiles. To adjust for potential confounding factors, three models were established as follows: model 1 adjusted for age and sex; model 2 adjusted for age, sex, BMI, smoking status, drinking status, physical activity, diabetes, dyslipidaemia, baPWV at baseline, and MAP at baseline and last follow-up; model 3 further adjusted for FBG (HDL-C if TG/HDL-C ratio analysed), triglyceride, PBG, LDL-C, eGFR, uric acid, homocysteine, and use of antidiabetic, lipid-lowering, or antihypertensive medications at baseline and follow-up. The regression coefficient and its 95% confidence interval (CI) are presented. Spearman's correlation analyses of the TyG index and TG/

HDL-C ratio with other common cardiometabolic risk factors were performed. To identify the interaction of insulin resistance indexes and other variables, the interactive terms were tested in the model.

All of the analyses presented above were conducted using R software (version 3.6.3). The difference was considered statistically significant at two-sided $P < 0.05$.

Results

The final analysis included 1,895 individuals. During the follow-up period, a total of 986 recruited individuals attended health examinations twice, 612 had three examinations and 297 had four or five examinations. The mean age of the population was 61.90 ± 12.75 years, and 1,477 (77.9%) were men. At baseline, 1,013 (53.5%) participants were diagnosed with hypertension, and 882 (46.5%) were diagnosed with prehypertension, among which 335 participants progressed to hypertension at the last visit of follow-up. The median values of the baPWV change, the baPWV change rate and the slope of baPWV were 46 cm/s, 9 cm/s/year and 8.11 in the whole population, respectively, as shown in Table 1. The detailed characteristics according to the quartiles of the TyG index and TG/HDL-C ratio are presented in Additional file 1: Table S1 and Table S2.

We observed linear and positive associations of the TyG index and TG/HDL-C ratio with the baPWV change, baPWV change rate and slope of baPWV, as shown in Fig. 2. In the fully adjusted model (model 3), a one-unit increase in the TyG index was associated with a 149.6 cm/s increase in baPWV change, a 40.4 cm/s/year increase in the baPWV change rate, and a 34.9 cm/s increase in the baPWV slope. The average increases of baPWV change, change rate and slope, comparing participants in the highest quartile versus the lowest of the TyG index, were 129.5 cm/s (P for trend: < 0.001), 37.6 cm/s/year (P for trend: < 0.001), and 30.6 (P for trend: 0.002), respectively. Similarly, a one-unit increase in the TG/HDL-C ratio was associated with a 37.1 cm/s increase in baPWV change, a 9.8 cm/s/year increase in the baPWV change rate, and an 8.0 cm/s increase in the baPWV slope. The average increases in baPWV change, change rate and slope, comparing participants in the highest quartile versus the lowest quartile of the TG/HDL-C ratio, were 133.4 cm/s (P for trend: 0.002), 43.5 cm/s/year (P for trend: 0.001), and 33.5 (P for trend: 0.004), respectively. The detailed regression results are shown in Tables 2 and 3. The distributions of baPWV change, baPWV change rate and slope of baPWV among the quartile groups according to the TyG index are shown in Fig. 3A–C. The distributions of baPWV parameters among the quartile groups according to the TG/HDL-C ratio are shown in Fig. 3D–F. To identify the significant

interaction of the TyG index and TG/HDL-C ratio with other covariates, such as age, sex, BMI, diabetes, dyslipidaemia, kidney function and life habits, we tested all of

A

P V

been reported to be associated with artery stiffness in the general population in previous studies. However, evidence in prehypertensive and hypertensive populations is scarce, except for the study by Li et al. [20], which reported that the TyG index is associated with evaluated baPWV in a hypertensive population based on a cross-sectional design. Our study supplemented the evidence that the TyG index and TG/HDL-C ratio are associated with artery stiffness progression for the first time. We found that a higher TyG index and TG/HDL-C ratio were associated with an increased baPWV change, baPWV change rate and slope of baPWV in the hypertensive population based on a prospective cohort design.

The TyG index and TG/HDL-C ratio, have been of increasing interest at present. The TyG index and TG/HDL-C ratio have high sensitivity and specificity for the diagnosis of insulin resistance compared with the hyperinsulinaemic-euglycaemic clamp and homeostatic model assessment in many populations [32, 33]. These two indicators have a high possibility of being easily applied in clinical practice for the early detection of IR, arterial stiffness and other diseases. Previous studies have

also demonstrated the association of the TyG index with cardiovascular, cerebrovascular and other metabolic diseases. In 2014, Fedchuk et al. [34] measured five steatosis biomarkers, including the TyG index, fatty liver index (FLI), NAFLD liver fat score (NAFLD-LFS), hepatic steatosis index (HSI) and visceral adiposity index (VAI), and found that all five steatosis biomarkers could contribute to the early diagnosis of steatosis and were correlated with IR. In the Northern Shanghai Study, Zhao et al. [17] concluded that an elevated TyG index was significantly related to arterial stiffness and nephric microvascular damage, which supported the clinical application of the TyG index for the assessment of vascular damage. In recent years, several studies have also reported that the TyG index may predict several cardiovascular diseases, including acute coronary syndromes, symptomatic CHD and ischaemic stroke [35–39]. For the TG/HDL-C ratio, in an 8-year Japan Diabetes Complications Study, Hirohito Sone et al. [40] evaluated conventional lipid variables, such as TG, non-HDL-C, TC/HDL-C ratio, LDL-C/HDL-C ratio, and TG/HDL-C ratio, for a relationship with CHD. According to their analyses, all

Table 2 Association of the TyG index with the absolute change, change rate and slope of baPWV


	model 1			model 2			model 3		
		95% CI	P value		95% CI	P value		95% CI	P value
Change of baPWV (cm/s)									
TyG (continuous)	32.034	4.642–59.425	0.022	57.119	22.835–91.402	0.001	149.582	90.823–208.342	<0.001
Quartile 2 (ref: quartile 1)	59.745	10.509–108.982	0.017	61.086	12.582–109.59	0.014	76.297	28.63–123.964	0.002
Quartile 3	85.613	36.416–134.81	0.001	93.631	43.889–143.373	<0.001	102.562	52.08–153.044	<0.001
Quartile 4	74.957	25.494–124.42	0.003	98.555	37.584–159.526	0.002	129.525	58.723–200.326	<0.001
P for trend			0.002			<0.001			<0.001
Change rate of baPWV (cm/s/year)									
TyG (continuous)	11.892	3.415–20.368	0.006	17.974	7.307–28.642	0.001	40.372	21.853–58.891	<0.001
Quartile 2 (ref: quartile 1)	11.870	-3.385–27.125	0.127	12.122	-2.986–27.229	0.116	16.529	1.507–31.551	0.031
Quartile 3	20.615	5.373–35.858	0.008	22.642	7.149–38.135	0.004	24.390	8.481–40.299	0.003
Quartile 4	24.616	9.291–39.941	0.002	30.750	11.76–49.741	0.002	37.607	15.294–59.919	0.001
P for trend			0.001			0.001			<0.001
Slope of baPWV									
TyG (continuous)	9.570	1.538–17.601	0.020	14.238	4.121–24.355	0.006	34.874	17.238–52.51	<0.001
Quartile 2 (ref: quartile 1)	10.986	-3.469–25.441	0.136	10.888	-3.439–25.215	0.137	14.872	0.57–29.175	0.042
Quartile 3	18.739	4.296–33.182	0.011	20.092	5.399–34.785	0.007	22.163	7.016–37.31	0.004
Quartile 4	20.126	5.604–34.647	0.007	23.648	5.639–41.658	0.010	30.576	9.332–51.82	0.005
P for trend			0.004			0.003			0.002

Model 1: Adjusted age and sex; Model 2: Age, sex, BMI, smoking status, drinking status, physical activity, diabetes, dyslipidaemia, baPWV at baseline, and MAP at baseline and follow-up; Model 3: Model 2 plus FBG, triglyceride, PBG, LDL-C, eGFR, uric acid, homocysteine, and use of antidiabetic, lipid-lowering, or antihypertensive medications at baseline and follow-up

Table 3 Association of TG/HDL-C ratio with the absolute change, change rate and slope of baPWV

	Model 1			Model 2			Model 3		
		95% CI	P value		95% CI	P value		95% CI	P value
Change of baPWV (cm/s)									
TG/HDL-C ratio (continuous)	6.795	-1.193 to 14.784	0.096	4.934	-3.684 to 13.552	0.062	37.057	7.85–66.264	0.013
Quartile 2 (ref: quartile 1)	70.320	21.037–119.604	0.005	68.883	19.964–117.802	0.006	75.317	24.101–126.534	0.004
Quartile 3	83.153	34.299–132.007	0.001	89.453	38.544–140.362	0.001	84.036	25.516–142.556	0.005
Quartile 4	131.912	82.542–181.282	<0.001	142.938	78.525–207.35	<0.001	133.420	51.949–214.891	0.001
P for trend			<0.001			<0.001			0.002
Change rate of baPWV (cm/s/year)									
TG/HDL-C ratio (continuous)	2.587	0.115–5.06	0.040	1.880	-0.801 to 4.562	0.169	9.768	0.547–18.989	0.038
Quartile 2 (ref: quartile 1)	14.121	-1.149 to 29.392	0.070	13.621	-1.616 to 28.857	0.080	19.486	3.325–35.646	0.018
Quartile 3	23.979	8.842–39.117	0.002	25.543	9.687–41.399	0.002	29.615	11.15–48.08	0.002
Quartile 4	38.237	22.94–53.534	<0.001	40.369	20.307–60.431	<0.001	43.511	17.804–69.217	0.001
P for trend			<0.001			<0.001			0.001
Slope of baPWV									
TG/HDL-C ratio (continuous)	2.253	-0.089 to 4.595	0.060	4.507	1.425–6.048	0.045	7.973	0.801–22.317	0.035
Quartile 2 (ref: quartile 1)	12.212	-2.264 to 26.687	0.098	11.300	-3.154 to 25.754	0.126	15.465	0.078–30.852	0.049
Quartile 3	21.636	7.287–35.986	0.003	21.977	6.935–37.019	0.004	24.296	6.715–41.877	0.007
Quartile 4	33.088	18.587–47.589	<0.001	32.178	13.146–51.21	0.001	33.496	9.02–57.972	0.007
P for trend			<0.001			<0.001			0.004

Model 1: adjusted age and sex; Model 2: age, sex, BMI, smoking status, drinking status, physical activity, diabetes, dyslipidaemia, baPWV at baseline, and MAP at baseline and follow-up; Model 3: Model 2 plus HDL-C, triglyceride, PBG, LDL-C, eGFR, uric acid, homocysteine, and use of antidiabetic, lipid-lowering, or antihypertensive medications at baseline and follow-up



Strengths of the present study include the prospective cohort design to explore the association of the TyG index and TG/HDL-C ratio with the progression of artery stiffness in prehypertensive and hypertensive populations, the adjustment of the potential confounding factors, and handling the TyG index and TG/HDL-C ratio as both continuous variables and categorical variables to enhance the reliability of our findings. However, the results should be interpreted in the context of some limitations. First, the sample size was relatively small, and the 95% CI of the estimated effect was wide. In the prehypertensive population, we only observed the evaluated tendency of baPWV change, baPWV change rate and slope of baPWV without statistical significance. The observed results in this single-centre study need further validation to generalize the associations in other populations and to evaluate the underlying biological mechanism between insulin resistance and artery stiffness progression. Second, the use of antihypertensive medication was considered in this analysis. However, we failed to collect the specific types of antihypertensive agents, given that

different antihypertensive medications could have different influences on baPWV measurements. Data on antiplatelet medication were unavailable. Third, we analysed the association of the TyG index and TG/HDL-C ratio at baseline with artery stiffness progression in the current study, and evaluation of the dynamic TyG index and TG/HDL-C ratio was needed with multiple measurements and related methods, such as trajectory analysis.

Conclusion

In summary, our findings indicate that the TyG index and TG/HDL-C ratio are significantly associated with arterial stiffness progression in hypertensive population, not in prehypertensive population. Monitoring the TyG index and TG/HDL-C ratio deserves more attention in clinical practice for the early prevention of arterial stiffness progression and other vascular complications of hypertension.

Abbreviations

TyG: Triglyceride glucose index; TG/HDL-C: Triglyceride to high-density lipoprotein cholesterol ratio; baPWV: Brachial-ankle pulse wave velocity; IR: Insulin resistance; BHMC: Beijing health management cohort.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12933-021-01330-6>.

Additional le 1: Table S1. Baseline characteristics by quartile groups of TyG index. **Table S2.** Baseline characteristics by quartiles of TG/HDL-C ratio. **Table S3.** The association of insulin resistance indexes and arterial stiffness progression in prehypertensive and hypertensive populations. **Table S4.** Joint relationship of TyG index and TG/HDL-C ratio with the absolute change, change rate and slope of baPWV. **Figure S1.** The correlation of TyG index and TG/HDL-C ratio with the cardio-metabolic risk factors.

Acknowledgements

We thank all the staff and participants of the Beijing health management cohort for their invaluable contributions.

Authors' contributions

Study conception and design: XG, WW, and LT; Data collection: XM, YL, JW, and ZH; Data analysis and interpretation: ZW, XL, ZL, and XL; Manuscript writing and reviewing: ZW, and DZ; Study supervision: XG, and LT. All authors read and approved the final manuscript.

Funding

Our work was funded by the National Natural Science Foundation of China (numbers: 81872708 and 82073668 to LX. Tao).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author (Dr. Lixin Tao) on reasonable request.

Declarations

Ethics approval and consent to participate

The study was approved by the Ethics Committees of Capital Medical University. All participants gave informed consent to participate before taking part. The approval number was 2020SY031.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Beijing Municipal Key Laboratory of Clinical Epidemiology, Department of Epidemiology and Health Statistics, School of Public Health, Capital Medical University, Beijing, China. ²Department of Mathematics and Statistics, La Trobe University, Melbourne, Australia. ³School of Medical and Health Sciences, Edith Cowan University, Perth, Australia.

Received: 6 May 2021 Accepted: 28 June 2021

Published online: 06 July 2021

References

- Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, et al. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation*. 2016;134(6):441–50.
- Lu J, Lu Y, Wang X, Li X, Linderman GC, Wu C, et al. Prevalence, awareness, treatment, and control of hypertension in China: data from 1.7 million adults in a population-based screening study (China PEACE Million Persons Project). *Lancet*. 2017;390(10112):2549–58.
- Badji A, Sabra D, Bherer L, Cohen-Adad J, Girouard H, Gauthier CJ. Arterial stiffness and brain integrity: A review of MRI findings. *Ageing Res Rev*. 2019;53:100907.
- Hughes TM, Wagenknecht LE, Craft S, Mintz A, Heiss G, Palta P, et al. Arterial stiffness and dementia pathology: Atherosclerosis Risk in Communities (ARIC)-PET Study. *Neurology*. 2018;90(14):e1248–56.
- Ahmed TAN, Shams-Eddin H, Fathy MA, El-Naggar HM, Kishk YT. Subclinical left ventricular systolic dysfunction by two-dimensional speckle-tracking echocardiography and its relation to ambulatory arterial stiffness index in hypertensive patients. *J Hypertens*. 2020;38(5):864–73.
- Aisu H, Saito M, Inaba S, Morofuji T, Takahashi K, Sumimoto T, et al. Association of worsening arterial stiffness with incident heart failure in asymptomatic patients with cardiovascular risk factors. *Hypertens Res*. 2017;40(2):173–80.
- Åström Malm I, Alehagen U, Blomstrand P, Dahlström U, De Basso R. Higher blood pressure in elderly hypertensive females, with increased arterial stiffness and blood pressure in females with the Fibrillin-1 2/3 genotype. *BMC Cardiovasc Disord*. 2020;20(1):180.
- Sun X, Chen Y, Zeng Q, Huang X, Cai J. Reduction of leukocyte-derived H(2)S linked to abnormal glycolipid metabolism in hypertensive subjects. *Clin Exp Hypertens*. 2017;39(5):427–34.
- Muniyappa R, Lee S, Chen H, Quon MJ. Current approaches for assessing insulin sensitivity and resistance in vivo: advantages, limitations, and appropriate usage. *Am J Physiol Endocrinol Metab*. 2008;294(1):E15–26.
- Guerrero-Romero F, Simental-Mendía LE, González-Ortiz M, et al. The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp. *J Clin Endocrinol Metab*. 2010;95(7):3347–51.
- Zhang M, et al. Cumulative increased risk of incident type 2 diabetes mellitus with increasing triglyceride glucose index in normal-weight people: The Rural Chinese Cohort Study. *Cardiovasc Diabetol*. 2017;16(1):30. <https://doi.org/10.1186/s12933-017-0514-x>.
- Florentino TV, Marini MA, Succurro E, Andreozzi F, Sesti G. Relationships of surrogate indexes of insulin resistance with insulin sensitivity assessed by euglycemic hyperinsulinemic clamp and subclinical vascular damage. *BMJ Open Diabetes Research Care*. 2019;7(1):e000911.
- Wu Z, Wang J, Li Z, Han Z, Miao X, Liu X, Li X, Wang W, Guo X, Tao L. Triglyceride glucose index and carotid atherosclerosis incidence in the Chinese population: A prospective cohort study. *Nutri Metab Cardiovasc Dis*. 2021. <https://doi.org/10.1016/j.numecd.2021.03.027>.
- Giannini C, Santoro N, Caprio S, Kim G, Lartaud D, Shaw M, et al. The triglyceride-to-HDL cholesterol ratio: association with insulin resistance in obese youths of different ethnic backgrounds. *Diabetes Care*. 2011;34(8):1869–74.
- Lee SB, Ahn CW, Lee BK, Kang S, Nam JS, You JH, et al. Association between triglyceride glucose index and arterial stiffness in Korean adults. *Cardiovasc Diabetol*. 2018;17(1):41.
- Won KB, Park EJ, Han D, Lee JH, Choi SY, Chun EJ, et al. Triglyceride glucose index is an independent predictor for the progression of coronary artery calcification in the absence of heavy coronary artery calcification at baseline. *Cardiovasc Diabetol*. 2020;19(1):34.
- Zhao S, Yu S, Chi C, Fan X, Tang J, Ji H, et al. Association between macro- and microvascular damage and the triglyceride glucose index in community-dwelling elderly individuals: the Northern Shanghai Study. *Cardiovasc Diabetol*. 2019;18(1):95.
- Won KB, Park GM, Lee SE, Cho JJ, Kim HC, Lee BK, et al. Relationship of insulin resistance estimated by triglyceride glucose index to arterial stiffness. *Lipids Health Dis*. 2018;17(1):268.
- Irace C, Carallo C, Scavelli FB, De Franceschi MS, Esposito T, Tripolino C, et al. Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index. *Int J Clin Pract*. 2013;67(7):665–72.
- Li M, Zhan A, Huang X, Hu L, Zhou W, Wang T, et al. Positive association between triglyceride glucose index and arterial stiffness in hypertensive patients: the China H-type Hypertension Registry Study. *Cardiovasc Diabetol*. 2020;19(1):139.

21. Liu J, Zhao Z, Mu Y, Zou X, Zou D, Zhang J, et al. Gender differences in the association between serum uric acid and prediabetes: a six-year longitudinal cohort study. *Int J Environ Res Public Health*. 2018;15(7):1560.
22. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT, et al. The seventh report of the joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure The JNC 7 report. *JAMA*. 2003;289(19):2560–72. <https://doi.org/10.1001/jama.289.19.2560>.
23. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150:604–12.
- 24.

Read your book online and receive BMC articles for free:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

